

Table 1. Clinical and demographic characteristics of PiB-positive and PiB-negative MCI subjects.

Characteristics	PiB-positive MCI	PiB-negative MCI	<i>p</i> -value
Sex (M:F)	4 : 8	6 : 5	0.30
Age (years)	74.0 ± 5.5	74.4 ± 10.8	0.58
Education (years)	11.2 ± 1.6	12.6 ± 1.2	0.04
MMSE	25.7 ± 1.8	25.4 ± 2.5	0.76
ADAS-Jcog	9.9 ± 2.5	9.3 ± 6.1	0.44
mean CBF	39.7 ± 2.7	38.8 ± 4.0	0.46
mean SUVR	1.96 ± 0.37	0.87 ± 0.2	< 0.0001

Data are shown as mean ± SD, MMSE: Mini-Mental State Examination, ADAS-Jcog: Alzheimer's Disease Assessment Scale-cognitive component- Japanese version, SUVR: standardized uptake value ratio of PiB, CBF: cerebral blood flow.

The  $\chi^2$  test for sex distribution and the Mann-Whitney U-test for age at examination, MMSE, ADAS-Jcog, mean SUVR values, and mean CBF were used to compare the characteristics of the two study subgroups. A *p*-value of less than 0.05 was considered statistically significant.